



Time left 0:01:47

## Question 19

Not yet answered

Marked out of 2.50

Flag question

If  $A = \begin{pmatrix} -1 & 4 \\ 3 & 1 \end{pmatrix}$ ,  $B = \begin{pmatrix} 2 & 3 \\ 1 & 0 \end{pmatrix}$ , and  $C = B^T \times A$ , then  $c_{21} =$


- ☐ A. 1
- ☐ B. -1
- ☐ C. 9
- ☐ D. 12
- ☒ E. -3

[Clear my choice](#)[Next page](#)

## Question 2

Not yet answered

Marked out of 2.50

 Flag question

Let  $A$  and  $B$  be two  $7 \times 7$  matrices such that  $\text{rk}(A) = 2$  and  $\text{rk}(B) = 3$ . Which of the following COULD be true?

- I.  $\text{rk}(A + B) = 0$
- II.  $\text{rk}(A + B) = 1$
- III.  $\text{rk}(A + B) = 6$

- ☒ A. I and II
- ☐ B. I, II and III
- ☐ C. I only
- ☐ D. None
- ☐ E. II only

[Clear my choice](#)



Time left 0:49:29

## Question 2

Not yet answered

Marked out of 2.50

Flag question

Let  $A$  and  $B$  be two  $7 \times 7$  matrices such that  $\text{rk}(A) = 2$  and  $\text{rk}(B) = 3$ . Which of the following COULD be true?

- I.  $\text{rk}(A + B) = 0$
- II.  $\text{rk}(A + B) = 1$
- III.  $\text{rk}(A + B) = 6$

- ☒ A. I and II
- ☐ B. I, II and III
- ☐ C. I only
- ☐ D. None
- ☐ E. II only

[Clear my choice](#)

[Next page](#)

Previous activity

[◀ Midterm Exam](#)





## PRINCIPLES OF LINEAR ALGEBRA

General

Final Exam

Time left 0:57:34

## Question 1

Not yet answered

Marked out of 2.50

Flag question

Let  $A = \begin{bmatrix} 0 & 1 & 0 & 0 \end{bmatrix}$ ,  $A^{-1}$  is equal to \_\_\_\_\_

- ☐ A. None of the mentioned
- ☒ B. Does not exist
- ☐ C. Identity matrix
- ☐ D. Null matrix

[Clear my choice](#)[Next page](#)



Time left 0:46:27

Question 4

Not yet answered

Marked out of 2.50

Flag question

The dot product of two vectors  $\vec{A}$  and  $\vec{B}$   
 $\vec{A} = 3i + 5j + 7k$   
 $\vec{B} = 11i + 13j + 17k$   
most nearly is

- ☐ A. 33
- ☒ B. 217
- ☐ C. 14.8
- ☐ D. 56


[Clear my choice](#)

[Next page](#)

[Previous activity](#)

Not yet answered

Marked out of 2.50

 Flag question

Let  $\{a, b\}$  and  $\{c, d\}$  be two sets of vectors, both linearly independent. Which of the following MUST be true?

- I.  $\{a, b, c, d\}$  is also linearly independent.
- II. Both  $\{a, c\}$  and  $\{b, d\}$  are linearly independent
- III. Either  $\{a, c\}$  or  $\{b, d\}$  is linearly independent

- ☐ A. III
- ☐ B. All are true
- ☐ C. II
- ☐ D. Neither is true
- ☒ E. I

[Clear my choice](#)[Next page](#)



## Question 5

Not yet answered

Marked out of 2.50

Flag question

Which of the following could be the set of eigenvalues of  $\begin{pmatrix} -2 & -4 & -4 \\ 2 & -2 & 2 \\ -2 & 4 & 0 \end{pmatrix}$ ?

- ☐ A.  $\lambda_1 = 2, \lambda_2 = 4, \lambda_3 = -2$
- ☒ B.  $\lambda_1 = 2, \lambda_2 = -4, \lambda_3 = -2$
- ☐ C.  $\lambda_1 = 2, \lambda_2 = 4, \lambda_3 = 2$
- ☐ D.  $\lambda_1 = 2, \lambda_2 = -4, \lambda_3 = 2$

[Clear my choice](#)



Time left 0:28:23

Question 9

Not yet answered

Marked out of 2.50

Flag question

The determinant  $\begin{vmatrix} -1 & -12 & -8 \\ 0 & 4 & 8 \\ 0 & 8 & -8 \end{vmatrix}$  is equal to

- ☒ A. None of these
- ☐ B. 52
- ☐ C. 64
- ☐ D. 0
- ☐ E. 4

[Clear my choice](#)[Next page](#)[Previous activity](#)[◀ Midterm Exam](#)






Question 5

Time left 0:42:03

Not yet answered

Marked out of 2.50

 Flag question

Which of the following could be the set of eigenvalues of  $\begin{pmatrix} -2 & -4 & -4 \\ 2 & -2 & 2 \\ -2 & 4 & 0 \end{pmatrix}$ ?

- ☐ A.  $\lambda_1 = 2, \lambda_2 = 4, \lambda_3 = -2$
- ☒ B.  $\lambda_1 = 2, \lambda_2 = -4, \lambda_3 = -2$
- ☐ C.  $\lambda_1 = 2, \lambda_2 = 4, \lambda_3 = 2$
- ☐ D.  $\lambda_1 = 2, \lambda_2 = -4, \lambda_3 = 2$

[Clear my choice](#)[Next page](#)[Previous activity](#)[◀ Midterm Exam](#)[Jump to...](#)[Next activity](#)



Marked out of 2.50

Time left 0:33:44

Flag question

The set of equations

$$\begin{bmatrix} 1 & 2 & 5 \\ 2 & 3 & 7 \\ 5 & 8 & 19 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 18 \\ 26 \\ 70 \end{bmatrix}$$

has

- ☐ A. No solution
- ☐ B. A unique solution
- ☐ C. Finite number of solutions
- ☒ D. Infinite solutions

[Clear my choice](#)[Next page](#)[Previous activity](#)[◀ Midterm Exam](#)



Time left 0:38:19

Question 6

Not yet answered

Marked out of 2.50

Flag question

Let  $A$  be the matrix that is inverse to  $\begin{pmatrix} 6 & -5 & -2 \\ -1 & 0 & -1 \\ -2 & 2 & 1 \end{pmatrix}$ . Then  $a_{22} =$

- ☐ A. 1
- ☐ B. 5
- ☐ C. None of these
- ☐ D. 3
- ☒ E. 2

[Clear my choice](#)

[Next page](#)

[Previous activity](#)

[◀ Midterm Exam](#)





Question 8

Time left 0:30:40

Not yet answered

Marked out of 2.50

Flag question

Which, if any, of these matrices have an  $LU$  decomposition?

☒ A. 
$$\begin{bmatrix} 1 & -3 & 7 \\ -2 & 6 & 1 \\ 0 & 3 & -2 \end{bmatrix}.$$

☐ B. None

☐ C. 
$$\begin{bmatrix} 3 & 2 \\ 0 & 1 \end{bmatrix}$$

☐ D. 
$$\begin{bmatrix} 0 & 1 \\ 3 & 2 \end{bmatrix}$$

[Clear my choice](#)[Next page](#)



Question 11

Time left 0:21:14

Not yet answered

Marked out of 2.50

Flag question

Let  $A$  be a  $3 \times 3$  matrix and let  $B = \begin{pmatrix} 5 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$ . If  $A$  is multiplied by  $B$  from the left then

- ☐ A. The 1st column of  $A$  is multiplied by 5
- ☒ B. None of these
- ☐ C. The 1st column of  $A$  is divided by 5
- ☐ D. The 1st row of  $A$  is multiplied by 5
- ☐ E. The 1st row of  $A$  is divided by 5

[Clear my choice](#)[Next page](#)[Previous activity](#)[◀ Midterm Exam](#)



General

Final Exam

Time left 0:23:43

Question 10

Not yet answered

Marked out of 2.50

Flag question

The eigenvalues of a  $4 \times 4$  matrix  $[A]$  are given as 2, -3, 13, and 7. The  $|\det(A)|$  then is

- ☐ A. 546
- ☒ B. 19
- ☐ C. Can not be determined
- ☐ D. 25

[Clear my choice](#)[Next page](#)[Previous activity](#)[Midterm Exam](#)





Time left 0:13:58

## Question 12

Not yet answered

Marked out of 2.50

Flag question

Let  $A$ ,  $B$ , and  $C$  be square matrices such that  $A \cdot B = E$  and  $B \cdot C = E$ , where  $E$  is the identity matrix. Which of the following MUST be true?

- I.  $\det(A) = \det(C)$
- II.  $A = C$
- III. If  $A = B$  then  $|\det(A)| = 1$

- ☐ A. I
- ☐ B. I and II
- ☐ C. II
- ☒ D. I, II and III
- ☐ E. III

Clear my choice



Time left 0:07:37

## Question 15

Not yet answered

Marked out of 2.50

Flag question

Consider there are only two computer companies in a country. The companies are named *Dude* and *Imac*. Each year, company *Dude* keeps  $1/5^{\text{th}}$  of its customers, while the rest switch to *Imac*. Each year, *Imac* keeps  $1/3^{\text{rd}}$  of its customers, while the rest switch to *Dude*. If in 2003, *Dude* had  $1/6^{\text{th}}$  of the market and *Imac* had  $5/6^{\text{th}}$  of the market, what will be share of *Dude* computers when the market becomes stable?

- ☐ A.  $6/11$
- ☐ B.  $53/90$
- ☐ C.  $37/90$
- ☒ D.  $5/11$

[Clear my choice](#)[Next page](#)[Previous activity](#)[◀ Midterm Exam](#)



Time left 0:11:23

Question **13**

Not yet answered

Marked out of 2.50

Flag question

Let  $A$  be a  $3 \times 4$  matrix and  $B$  be a  $4 \times 5$  matrix.  $\text{rk}(A \cdot B)$  is not greater than

- ☐ A. Could be any number
- ☐ B. 4
- ☐ C. 5
- ☒ D. 3
- ☐ E. 0

[Clear my choice](#)

[Next page](#)

[Previous activity](#)

[Midterm Exam](#)





Time left 0:08:33

Question 14

Not yet answered

Marked out of 2.50

Flag question

If  $\begin{bmatrix} -4.5 \\ -4 \\ 1 \end{bmatrix}$  is an eigenvector of  $\begin{bmatrix} 8 & -4 & 2 \\ 4 & 0 & 2 \\ 0 & -2 & -4 \end{bmatrix}$ , the eigenvalue corresponding to the eigenvector is

- ☒ A. 4
- ☐ B. 1
- ☐ C. -4.5
- ☐ D. 6

[Clear my choice](#)[Next page](#)[Previous activity](#)[◀ Midterm Exam](#)

Jump to





Time left 0:04:23

The lower triangular matrix  $[L]$  in the LU decomposition of the matrix

$$\begin{bmatrix} 25 & 5 & 4 \\ 10 & 8 & 16 \\ 8 & 12 & 22 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ \ell_{21} & 1 & 0 \\ \ell_{31} & \ell_{32} & 1 \end{bmatrix} \begin{bmatrix} u_{11} & u_{12} & u_{13} \\ 0 & u_{22} & u_{23} \\ 0 & 0 & u_{33} \end{bmatrix}$$

is

☐ A. 
$$\begin{bmatrix} 1 & 0 & 0 \\ 0.40000 & 1 & 0 \\ 0.32000 & 1.5000 & 1 \end{bmatrix}$$

☐ B. 
$$\begin{bmatrix} 1 & 0 & 0 \\ 10 & 1 & 0 \\ 8 & 12 & 0 \end{bmatrix}$$

☐ C. 
$$\begin{bmatrix} 25 & 5 & 4 \\ 0 & 6 & 14.400 \\ 0 & 0 & -4.2400 \end{bmatrix}$$

☒ D. 
$$\begin{bmatrix} 1 & 0 & 0 \\ 0.40000 & 1 & 0 \\ 0.32000 & 1.7333 & 1 \end{bmatrix}$$


[Clear my choice](#)

## Question 18

Time left 0:03:12

Not yet answered

Marked out of 2.50

 Flag question

Let  $A$  be a  $3 \times 3$  matrix with eigenvalues  $\lambda_1$ ,  $\lambda_2$ , and  $\lambda_3$  such that  $\lambda_1 \neq \lambda_2$ ,  $\lambda_1 \neq \lambda_3$ , and  $\lambda_2 \neq \lambda_3$ . Which of the following MUST be true?

- I. If  $a$  and  $b$  are eigenvectors corresponding to  $\lambda_1$  then the set  $\{a, b\}$  is linearly dependent.
- II. If  $a_1$  and  $a_2$  are eigenvectors corresponding to  $\lambda_1$  and  $\lambda_2$ , respectively, then  $\{a_1, a_2\}$  is linearly independent.
- III.  $\det(A) \neq 0$ .

- ☐ A. I
- ☐ B. III
- ☐ C. I and II
- ☐ D. II
- ☐ E. II and III

Next page





Time left 0:06:50

## Question 16

Not yet answered

Marked out of 2.50

Flag question

Let  $A$  be a  $3 \times 3$  matrix whose characteristic polynomial  $f(\lambda) = 1 + \lambda - \lambda^2 - \lambda^3$ . Which of the following MUST be true?

- I.  $\text{rk}(A - \lambda I) = 3$  for some  $\lambda$
- II.  $A^{-1}$  exists
- III.  $A$  is NOT diagonalizable

- ☐ A. II
- ☐ B. I
- ☐ C. I, II, and III
- ☐ D. I and II
- ☒ E. II and III

[Clear my choice](#)[Next page](#)[Previous activity](#)[◀ Midterm Exam](#)